

Note on f_λ -statistical convergence

Stuti Borgohain ^{*1} and Ekrem Savaş

Istanbul Commerce University, 34840 Istanbul, Turkey

E-mail : stutiborgohain@yahoo.com, ekremsavas@yahoo.com.

Abstract: In this article, we study about the λ -statistical convergence with respect to the density of moduli and find some results related to statistical convergence as well. Also we introduce the concept of f_λ -summable sequence and try to investigate some relation between the f_λ -summability and module λ -statistical convergence.

1 Introduction

The idea of statistical convergence of real sequences is the extension of convergence of real sequences. Initially the statistical convergence for single sequences was introduced by Fast [18] in 1951 and Schoenberg [23] in 1959, studied statistical convergence as a summability method and listed some of elementary properties of statistical convergence as well. Both of these authors noted that if bounded sequence is statistically convergent, then it is Cesaro summable. More investigations have been studied in the direction of topological spaces, statistical Cauchy condition in uniform spaces, Fourier analysis, ergodic theory, number theory, measure theory, trigonometric series, turnpike theory and Banach spaces. Statistical convergence turned out to be one of the most active areas of research in summability theory after the works of Fridy [25] and Salat [28]. Recently, Savas [5] introduced the generalized double statistical convergence in locally solid Riesz spaces. Moreover, Mursaleen [26] introduced the concept of λ -statistically convergence by using the idea of (V, λ) -summability to generalize the concept of statistical convergence. For more interesting investigations concerning statistical convergence one may consult the papers of Cakalli [16], Miller and Orhan [17]

¹The work of the first author was carried during her one month visit to Istanbul Ticaret University, Turkey, in November'2015

and ([24], [3], [4], [27], [28], [13], [2], [?]) and others.

The concept of density for sets of natural numbers with respect to the modulus function was introduced by A. Aizpuru et al [1] in 2014. They studied and characterized the generalization of this notion of f -density with statistical convergence and proved that ordinary convergence is equivalent to the module statistical convergence for every unbounded modulus function. Also, in [2], A. Aizpuru and his team, they worked on double sequence spaces for the results of f -statistical convergence by using unbounded modulus function. The concept was further generalized and characterized by Savas and Borgohain [[?], [6]] .

The notion of statistical convergence depends on the idea of asymptotic density of subsets of the set $\mathbb{N}$ of natural numbers. A subset A of $\mathbb{N}$ is said to have natural density $\delta(A)$ if

$$\delta(A) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \chi_A(k).$$

where χ_A is the characteristic function of A .

We mean a sequence (x_t) to be statistically convergent to L , if for any $\xi > 0$, $\delta(\{t \in \mathbb{N} : |x_t - L| \geq \xi\}) = 0$. Analogously, (x_t) is said to be statistically Cauchy if for each $\xi > 0$ and $n_0 \in \mathbb{N}$ there exists an integer $q \geq n_0$ such that $\delta(\{t \in \mathbb{N} : \|x_t - x_q\| < \xi\}) = 1$.

Nakano [19] introduced the notion of a modulus function whereas Ruckle [29] and Maddox [21] have introduced and discussed some properties of sequence spaces defined by using a modulus function. A modulus function is a function $f : R^+ \rightarrow R^+$ which satisfies:

1. $f(x) = 0$ if and only if $x = 0$.
2. $f(x + y) \leq f(x) + f(y)$ for every $x, y \in R^+$.
3. f is increasing.
4. f is continuous from the right at 0.

Later on modulus function have been discussed in ([20, 10, 11, 12, 14]) and others.

In this paper, we study the density on moduli with respect to the λ -statistical convergence. We also investigate some results on the new concept of f_λ -statistical convergence with the ordinary convergence. Also we find out some new concepts on f_λ -summability theory and try to find out new results related to the f_λ -summable sequences and f_λ -statistical convergent sequence.

2 Definitions and basic results

By density of moduli of a set $A \subseteq \mathbb{N}$, we mean $\delta_f(A) = \lim_u \frac{f(|A(u)|)}{f(u)}$, (in case this limit exists) where f is an unbounded modulus function.

Let (x_t) be a sequence in X (X is a normed space). If for each $\xi > 0$, $A = \{t \in \mathbb{N} : \|x_t - L\| > \xi\}$ has f -density zero, then it is said that the f -statistical limit of (x_t) is $L \in X$, and we write it as $f\text{-stlim} x_t = L$.

Observe that $\delta(A) = 1 - \delta(\mathbb{N} \setminus A)$.

Let us assume that $A \subseteq \mathbb{N}$ and A has f -density zero. For every $u \in \mathbb{N}$ we have $f(u) \leq f(|A(u)|) + f(|(\mathbb{N} \setminus A)(u)|)$ and so

$$1 \leq \frac{f(|A(u)|)}{f(u)} + \frac{f(|(\mathbb{N} \setminus A)(u)|)}{f(u)} \leq \frac{f(|A(u)|)}{f(u)} + 1$$

By taking limits we deduce that f -density of $(\mathbb{N} \setminus A)$ is one.

Let $\lambda = (\lambda_n)$ be a non-decreasing sequence of positive numbers such that ,

$$\lambda_1 = 1, \lambda_{n+1} \leq \lambda_n + 1 \text{ and } \lambda_n \rightarrow \infty \text{ as } n \rightarrow \infty.$$

Note: The collection of all such sequences λ will be denoted by Λ .

A sequence (x_t) of real numbers is said to be λ -statistically convergent to L if for any $\xi > 0$,

$$\lim_{n \rightarrow \infty} \frac{1}{\lambda_n} |\{t \in I_n : |x_t - L| \geq \xi\}| = 0,$$

where $I_n = [n - \lambda_n + 1, n]$ and $|A|$ denotes the cardinality of $A \subset N$ (refer [26]).

More investigations in this direction and more applications of ideals can be found in [5, 8, 9, 15] where many important references can be found.

We define f_λ -statistical convergence as:

$$\lim_{n \rightarrow \infty} \frac{f(|A(n)|)}{f(\lambda_n)} = 0 \text{ i.e. } \delta_{f_\lambda}(A) = \lim_{n \rightarrow \infty} \frac{f(|A(n)|)}{f(\lambda_n)}.$$

The set of all f_λ -statistically convergent sequences is denoted by S_{f_λ} .

A sequence $x = (x_t)$ is said to be strongly f_λ -summable to the limit L if

$$\lim_n \frac{1}{f(\lambda_n)} \sum_{t \in I_n} f(|x_t - L|) = 0$$

and we write it as $x_t \rightarrow L[f_\lambda]$. In this case, L is called the f_λ -limit of x . We denote the class of all f_λ -summable sequences as w_{f_λ} .

3 Main Results

Maddox [22] showed the existence of an unbounded modulus f for which there exists a positive constant c such that $f(xy) \geq cf(x)f(y)$ for all $x \geq 0, y \geq 0$.

Theorem 3.1. *Let f be an unbounded modulus such that there is a positive constant c such that $f(xy) \geq cf(x)f(y)$ for all $x \geq 0, y \geq 0$ and $\lim_{u \rightarrow \infty} \frac{f(u)}{u} > 0$. A strongly λ -summable sequence (x_t) is also a f_λ -statistically convergent sequence. Moreover, for a bounded sequence (x_t) , f_λ -statistical convergence implies strongly λ -summability.*

Proof. By the definition of modulus function, we have for any sequence $x = (x_t)$ and $\xi > 0$,

$$\begin{aligned} \sum_{t=1}^n f(|x_t - L|) &\geq f\left(\sum_{t=1}^n f(|x_t - L|)\right) \\ &\geq f(|\{t \leq n : |x_t - L| \geq \xi\}|\xi) \\ &\geq cf(|\{t \leq n : |x_t - L| \geq \xi\}|)f(\xi) \end{aligned}$$

Since $x = (x_t)$ is strongly λ -summable sequence, so

$$\begin{aligned} \frac{1}{\lambda_n} \sum_{t=1}^n f(|x_t - L|) &\geq \frac{cf(\{t \leq n : |x_t - L| \geq \xi\})f(\xi)}{\lambda_n} \\ &= \frac{cf(\{t \leq n : |x_t - L| \geq \xi\})f(\xi)f(\lambda_n)}{\lambda_n f(\lambda_n)} \end{aligned}$$

By using the fact that $\lim_{u \rightarrow \infty} \frac{f(u)}{u} > 0$ and x is λ -statistical w.r.t. f , it can be summarized that x is f_λ -statistical convergent and this completes the proof of the theorem. $\square$

Theorem 3.2. For a strongly λ -summable or statistically f_λ -convergent sequence $x = (x_t)$ to L , there is a convergent sequence y and a f_λ -statistically null sequence z such that y is convergent to L , $x = y + z$ and $\lim_n \frac{f(\{t \in I_n : z_t \neq 0\})}{f(\lambda_n)} = 0$. Moreover, if x is bounded, then y and z both are bounded.

Note: Here f be an unbounded modulus such that there is a positive constant c such that $f(xy) \geq cf(x)f(y)$ for all $x \geq 0, y \geq 0$ and $\lim_{u \rightarrow \infty} \frac{f(u)}{u} > 0$.

Proof. From the previous theorem, we have x is strongly λ -summable to L implies x is f_λ -statistically convergent to L . Choose a strictly increasing sequence of positive integers $M_1 < M_2 < M_3 \dots$ such that,

$$\frac{1}{f(\lambda_n)} f\left(\left|\left\{t \in I_n : |x_t - L| \geq \frac{1}{d}\right\}\right|\right) < \frac{1}{d}, \text{ for } n > N_d, \text{ where } N_0 = 0$$

If $N_0 < t < N_1$, let us set $z_t = 0$ and $y_t = x_t$.

Let $d \geq 1$ and $N_d < t \leq N_{d+1}$, we set,

$$y_t = \begin{cases} x_t, z_t = 0 & \text{if } |x_t - L| < \frac{1}{d}; \\ L, z_t = x_t - L & \text{if } |x_t - L| \geq \frac{1}{d}. \end{cases}$$

Clearly, $x = y + z$ and y and z are bounded, if x is bounded.

Observe that $|y_t - L| < \xi$ for $t > N_d$ and $|y_t - L| = |x_t - L| < \xi$ if $|x_t - L| < \frac{1}{d}$.

Which follows that $|y_t - L| = |L - L| = 0$ if $|x_t - L| > \frac{1}{d}$.

Hence, for ξ arbitrary, we get $\lim_t y_t = L$.

Next, we observe that, $f(\{|t \in I_n : z_t \neq 0\}|) \geq f(\{|t \in I_n : |z_t| \geq \xi\}|)$, for any natural number n and $\xi > 0$. Hence,

$$\lim_n \frac{f(\{|t \in I_n : z_t \neq 0\}|)}{f(\lambda_n)} = 0;$$

that is z is f_λ -statistically null.

We now show that if $\beta > 0$ and $d \in \mathbb{N}$ such that $\frac{1}{d} < \beta$, then $f(\{|t \in I_n : z_t \neq 0\}|) < \beta$ for all $n > N_d$.

If $N_d < t \leq N_{d+1}$, then $z_t \neq 0$ only if $|x_t - L| > \frac{1}{d}$. It follows that if $N_p < t \leq N_{p+1}$, then,

$$\{t \in I_n : z_t \neq 0\} \subseteq \left\{t \in I_n : |x_t - L| > \frac{1}{p}\right\}.$$

Consequently,

$$\frac{1}{f(\lambda_n)} f(\{|t \in I_n : z_t \neq 0\}|) \leq \frac{1}{f(\lambda_n)} f\left(\left|\left\{t \in I_n : |x_t - L| > \frac{1}{p}\right\}\right|\right) < \frac{1}{p} < \frac{1}{d} < \beta,$$

if $N_p < n \leq N_{p+1}$ and $p > d$.

That is, $\lim_n \frac{1}{f(\lambda_n)} f(\{|t \in I_n : z_t \neq 0\}|) = 0$.

This completes the proof of the theorem.

□

Corollary 3.1. *Let f, g be unbounded moduli, X a normed space, (x_t) a sequence in X and $x, y \in X$. We have*

1. *A f_λ -statistical convergent sequence is also statistically convergent to the same limit.*

2. The f_λ -statistical limit is unique whenever it exists.

3. Let f_λ -statlim $x_t = x$ and f_λ -statlim $y_t = y$, then,

- f_λ -statlim $(x_t \pm y_t) = x \pm y$
- f_λ -statlim $\alpha x_t = \alpha x, \alpha \in \mathbb{R}$.

4. If f -stlim $x_t = x$ and g -stlim $x_t = y$ then $x = y$, i.e two different methods of statistical convergence are always compatible.

Theorem 3.3. Let (x_t) be a sequence in a normed space X . For an unbounded modulus f , (x_t) is f_λ -statistically convergent to x if and only if there exists $T \subseteq \mathbb{N}$ such that f_λ -density of T is zero and (x_t) has λ -summable to x .

Proof. Let us assume that $V_z = \left\{ t \in \mathbb{N} : \sum_t \|x_t - x\| > \frac{1}{z} \right\}$, for every $z \in \mathbb{N}$ such that $V_z \subset V_{z+1}$ and $\lim_n \frac{f(|V_z(n)|)}{f(\lambda_n)} = 0$.

Also by induction, we obtain $\frac{f(|V_z(n)|)}{f(\lambda_n)} \leq \frac{1}{z}$ whenever $n \geq i_z$.

Note: Here $i_1 < i_2 < i_3 \dots$ is a strictly increasing sequence of positive numbers such that $i_z \in V_z$.

Again set $T = \cup_{z \in \mathbb{N}} ([i_z, i_{z+1}) \cap V_z)$. Then for every $n \geq i_1$ there exists $z \in \mathbb{N}$ such that $i_z \leq n \leq i_{z+1}$ and if $m \in T(n)$ then $m < i_{z+1}$, which implies $m \in V_z$. Therefore $T(n) \subseteq V_z(n)$ and thus,

$$\frac{f(|T(n)|)}{f(\lambda_n)} \leq \frac{f(|V_z(n)|)}{f(\lambda_n)} \leq \frac{1}{z}$$

which follows that T has f_λ -density zero.

For $\xi > 0$ and $z \in \mathbb{N}$ such that $\frac{1}{z} < \xi$, we have $n \in \mathbb{N} \setminus T$ and $n \geq i_z$ for which there exists $q \geq z$ with $i_q \leq n \leq i_{q+1}$ and this implies $n \notin V_q$, so,

$$\frac{1}{\lambda_n} \sum_t \|x_t - x\| \leq \frac{1}{q} \leq \frac{1}{z} < \xi.$$

which shows that (x_t) has λ -summable to x .

Conversely, let us assume that for $T \in \mathbb{N}$, $\lim_{t \in \mathbb{N} \setminus T} \frac{1}{\lambda_n} \|x_t - x\| = 0$ for $n \in I_n$ and T has f_λ -density zero. For $\xi > 0$, there exists $t_0 \in \mathbb{N}$ such that if $t > t_0$ and $t \in \mathbb{N} \setminus T$ then

$$\frac{1}{\lambda_n} \sum_t \|x_t - x\| \leq \xi.$$

This implies,

$$\left\{ t \in \mathbb{N} : \sum_t \|x_t - x\| > \xi \right\} \subseteq T \cup \{1, \dots, t\}$$

$$\text{and then } \delta_{f_\lambda} \left(\left\{ t \in \mathbb{N} : \sum_t \|x_t - x\| > \xi \right\} \right) = 0.$$

□

Definition 3.1. The sequence (x_t) is f_λ -statistically Cauchy if for every $\xi > 0$ there exists $Q \in \mathbb{N}$ such that $\delta_{f_\lambda} \left(\left\{ t \in \mathbb{N} : \sum_t \|x_t - x_Q\| > \xi \right\} \right) = 0$.

Corollary 3.2. A sequence is f_λ -statistically convergent implies that it is f_λ -statistically Cauchy. The converge is true if the space is complete and this result is a particular case of filter convergence.

Theorem 3.4. For the Banach space X where f be an unbounded modulus, f_λ -statistically Cauchy sequence is f_λ -statistically convergent sequence.

Proof. For every $k \in \mathbb{N}$, let x_{Q_k} be such that

$$\delta_{f_\lambda} \left(\left\{ t \in \mathbb{N} : \sum_t \|x_t - x_{Q_k}\| > \frac{1}{k} \right\} \right) = 0.$$

Consider the set $J_k = \cap_{i \leq k} B(x_{Q_i}, \frac{1}{i})$, then for each $k \in \mathbb{N}$ we have $\text{diam}(J_k) \leq \frac{2}{k}$

and $R_k = \{t \in \mathbb{N} : x_t \notin J_k\}$ so that $R_k = \cup_{i \leq k} \left\{ t \in \mathbb{N} : \sum_t \|x_t - x_{Q_i}\| > \frac{1}{i} \right\}$ which implies f_λ -density of R_k is zero.

Following the proof of the previous theorem, we get for $t \geq r_k$ then $\frac{f(|R_k(t)|)}{f(\lambda_t)} \leq \frac{1}{k}, r_k \in R_k$.

Considering $D = \cup_{k \in \mathbb{N}} ([r_k, r_{k+1}) \cap R_k)$, we get f_λ -density of D is zero.

Since $\cap_{k \in \mathbb{N}} J_k$ has exactly one element, say x , due to the completeness of X . We have to prove that $\lim_{t \in \mathbb{N} \setminus D} \frac{1}{\lambda_n} \|x_t - x\| = 0$.

For $\xi > 0$, let us choose $i \in \mathbb{N}$ such that $\frac{2}{i} < \xi$. If $t \geq r_j$ and $t \in \mathbb{N} \setminus D$ then there exists $k \geq i$ such that $r_k \leq t < r_{k+1}$ and then $t \notin R_k$, which implies $x_t \in J_k$ and thus,

$$\frac{1}{\lambda_n} \sum_t \|x_t - x\| \leq \frac{2}{k} \leq \frac{2}{i} < \xi.$$

This completes the proof of the theorem. □

Corollary 3.3. *A sequence which is f_λ -statistically convergent for each module f , is also convergent in ordinary sense, i.e. if $B \subseteq \mathbb{N}$ is infinite, then there exists an unbounded module f such that f_λ -density of B is one.*

Theorem 3.5. *Let (x_t) be a sequence in X . If for every unbounded modulus f there exists f_λ -stlim x_t then all these limits are the same $x \in X$ and (x_t) is also λ -summable to x in the norm topology.*

Proof. It is proved that for f, g two unbounded moduli, the f_λ -statistical limit is unique whenever it exists.

Let X a normed space, (x_t) a sequence in X and $P, Q \in X$ and if the uniqueness is false that $\lim x_t = P$, there exists $\xi > 0$ such that

$$B = \left\{ t \in \mathbb{N} : \sum_p \|x_t - P\| > \xi \right\}$$

is infinite.

Now, by choosing an unbounded modulus f which will satisfy $\delta_{f_\lambda}(B) = 1$, then this clearly contradicts the assumption that $f_\lambda\text{-stlim } x_t = x$, which completes the proof of the Theorem. $\square$

Theorem 3.6. *If $\lambda \in \Lambda$ with $\liminf_{n \rightarrow \infty} \frac{n}{f(\lambda_n)} > 0$, then f_λ -statistically convergent sequences are f -statistical convergent.*

Proof. For given $\xi > 0$, we have,

$$\{t \leq n : |x_t - L| \geq \xi\} \supset \{t \in I_n : |x_t - L| \geq \xi\}.$$

Therefore,

$$\begin{aligned} \frac{1}{f(\lambda_n)} f(|\{t \leq n : |x_t - L| \geq \xi\}|) &\geq \frac{1}{f(\lambda_n)} f(|\{t \in I_n : |x_t - L| \geq \xi\}|) \\ &\geq \frac{f(n)}{n} \frac{n}{f(\lambda_n)} \frac{1}{f(n)} f(|\{t \in I_n : |x_t - L| \geq \xi\}|) \end{aligned}$$

Taking the limit as $n \rightarrow \infty$ and using the fact that $\lim_n \frac{f(n)}{n} > 0$, we get (x_t) is f_λ -statistical convergent implies (x_t) is f -statistical convergent. $\square$

Theorem 3.7. *Let $\lambda \in \Lambda$ be such that $\lim_n \frac{n}{f(\lambda_n)} = 1$. Then, f -statistical convergence is f_λ -statistical convergence.*

Proof. Let $\beta > 0$ be given. Since $\lim_n \frac{n}{f(\lambda_n)} = 1$, we can choose $m \in \mathbb{N}$ such that, $\frac{f(n-\lambda_n+1)}{f(n)} < \frac{\beta}{2}$ for all $n \geq m$.

Now observe that,

$$\begin{aligned}
\frac{1}{f(n)}f(|\{t \leq n : |x_t - L| \geq \xi\}|) &= \frac{1}{f(n)}f(|\{t < n - \lambda_n + 1 : |x_t - L| \geq \xi\}|) \\
&+ \frac{1}{f(n)}f(|\{t \in I_n : |x_t - L| \geq \xi\}|) \\
&< \frac{f(n - \lambda_n + 1)}{f(n)} + \frac{1}{f(n)}f(|\{t \in I_n : |x_t - L| \geq \xi\}|) \\
&< \frac{\beta}{2} + \frac{1}{f(\lambda_n)}f(|\{t \in I_n : |x_t - L| \geq \xi\}|)
\end{aligned}$$

Taking limit as $n \rightarrow \infty$, we get, (x_t) is f -statistically convergent is f_λ -statistically convergent. $\square$

Theorem 3.8. Let $\lambda = (\lambda_n)$ and $\mu = (\mu_n)$ be two sequences in Λ such that $\lambda_n \leq \mu_n$ for all $n \in \mathbb{N}_{n_0}$.

1. If $\liminf_{n \rightarrow \infty} \frac{\lambda_n}{\mu_n} > 0$ then $S_{f_\mu} \subseteq S_{f_\lambda}$.
2. If $\lim_{n \rightarrow \infty} \frac{\mu_n}{\lambda_n} = 1$ then $S_{f_\lambda} \subseteq S_{f_\mu}$.

Proof. 1. Suppose that $\lambda_n \leq \mu_n$ for all $n \in \mathbb{N}_{n_0}$ which implies that $f(\lambda_n) \leq f(\mu_n)$.

Given $\liminf_{n \rightarrow \infty} \frac{\lambda_n}{\mu_n} > 0$, then with the property of modulus function, we get $\liminf_{n \rightarrow \infty} \frac{f(\lambda_n)}{f(\mu_n)} > 0$. Also, since $\lambda_n \leq \mu_n$ for all $n \in \mathbb{N}_{n_0}$, so $I_n \subset J_n$ where $I_n = [n - \lambda_n + 1, n]$ and $J_n = [n - \mu_n + 1, n]$.

Now, for $\xi > 0$, we can write,

$$\{t \in J_n : |x_t - L| \geq \xi\} \supset \{t \in I_n : |x_t - L| \geq \xi\}$$

and so,

$$\frac{1}{\mu_n}|\{t \in J_n : |x_t - L| \geq \xi\}| \geq \frac{\lambda_n}{\mu_n} \frac{1}{\lambda_n}|\{t \in I_n : |x_t - L| \geq \xi\}|, \text{ for all } n \in \mathbb{N}_{n_0}.$$

Now, by using the definition of modulus function and satisfying $\liminf_{n \rightarrow \infty} \frac{\lambda_n}{\mu_n} > 0$ and also taking limit as $n \rightarrow \infty$, we get $S_{f_\mu} \subseteq S_{f_\lambda}$. $\square$

Proof. 2. Let $S_{f_\lambda} - \lim x_t = x$ and $\lim_{n \rightarrow \infty} \frac{\mu_n}{\lambda_n} = 1$.

Since $I_n \subset J_n$, for $\xi > 0$, we write,

$$\begin{aligned}
\frac{1}{\mu_n} |\{t \in J_n : |x_t - L| \geq \xi\}| &= \frac{1}{\mu_n} |\{n - \mu_n + 1 \leq k \leq n - \lambda_n : |x_t - L| \geq \xi\}| + \frac{1}{\mu_n} |\{t \in I_n : |x_t - L| \geq \xi\}| \\
&\leq \frac{\mu_n - \lambda_n}{\mu_n} + \frac{1}{\mu_n} |\{t \in I_n : |x_t - L| \geq \xi\}| \\
&\leq \frac{\mu_n - \lambda_n}{\lambda_n} + \frac{1}{\mu_n} |\{t \in I_n : |x_t - L| \geq \xi\}| \\
&\leq \left(\frac{\mu_n}{\lambda_n} - 1 \right) + \frac{1}{\lambda_n} |\{t \in I_n : |x_t - L| \geq \xi\}|, \text{ for all } n \in \mathbb{N}_{n_0}.
\end{aligned}$$

Since $\lim_{n \rightarrow \infty} \frac{\mu_n}{\lambda_n} = 1$ and using the definition of modulus function, we can say that $x = (x_t)$ is S_{f_μ} -statistically convergent is S_{f_λ} -statistically convergent sequence, so $S_{f_\lambda} \subseteq S_{f_\mu}$. $\square$

Corollary 3.4. Let $\lambda = (\lambda_n)$ and $\mu = (\mu_n)$ be two sequences in Λ such that $\lambda_n \leq \mu_n$ for all $n \in \mathbb{N}_{n_0}$. If $\liminf_{n \rightarrow \infty} \frac{\lambda_n}{\mu_n} > 0$, then $w_{f_\mu} \subset w_{f_\lambda}$.

Theorem 3.9. Let $\lambda_n \leq \mu_n$ for all $n \in \mathbb{N}_{n_0}$, then, if $\liminf_{n \rightarrow \infty} \frac{\lambda_n}{\mu_n} > 0$, then a strongly w_{f_μ} -summable sequence is S_{f_λ} -statistically convergent sequence.

Proof. For any $\xi > 0$, we have,

$$\begin{aligned}
\sum_{t \in J_n} f(|x_t - L|) &= \sum_{t \in J_n, f(|x_t - L|) \geq \xi} f(|x_t - L|) + \sum_{t \in J_n, f(|x_t - L|) < \xi} f(|x_t - L|) \\
&\geq \sum_{t \in I_n, f(|x_t - L|) \geq \xi} f(|x_t - L|) + \sum_{t \in I_n, f(|x_t - L|) < \xi} f(|x_t - L|) \\
&\geq \sum_{t \in I_n, f(|x_t - L|) \geq \xi} f(|x_t - L|) \\
&\geq f \left(\sum_{t \in J_n, f(|x_t - L|) \geq \xi} |x_t - L| \right) \\
&\geq f(|\{t \in I_n : |x_t - L| \geq \xi\}| \cdot \xi) \\
&\geq c f(|\{t \in I_n : |x_t - L| \geq \xi\}|) \cdot f(\xi)
\end{aligned}$$

and so

$$\begin{aligned}\frac{1}{\mu_n} \sum_{t \in I_n} f(|x_t - L|) &\geq \frac{1}{\mu_n} f(|\{t \in I_n : |x_t - L| \geq \xi\}|) f(\xi) \\ &\geq \frac{\lambda_n}{\mu_n} \frac{1}{\lambda_n} f(|\{t \in I_n : |x_t - L| \geq \xi\}|) f(\xi)\end{aligned}$$

Since $\liminf_{n \rightarrow \infty} \frac{\lambda_n}{\mu_n} > 0$ and by using the definition of modulus function, we get, (x_t) is strongly f_λ -summable to L implies (x_t) is S_{f_λ} statistically convergent to L .

This completes the proof of the theorem. □

References

- [1] A. Aizpuru, M.C. Listán-García and F. Rambla-Barreno, Density by Moduli and Statistical Convergence, Quaestiones Mathematicae 2014: 1-6. <http://dx.doi.org/10.2989/16073606.2014.894682>
- [2] B.C. Tripathy and S. Borgohain, Statistically Convergent Difference Sequence spaces of Fuzzy Real numbers defined by Orlicz Function, Thai Journal of Mathematics, 11(2) (2013), 357-370.
- [3] E. Kolk, The statistical convergence in Banach spaces, Acta Et Commentationes Univ. Tartuensis, 928(1991), 41-52.
- [4] E. Kolk, Matrix summability of statistically convergent sequences, Analysis, 13(1993), 77-83.
- [5] E. Savaş, On generalize ddouble statistical convergence in locally solid Riesz spaces, Miskolc Mathematical Notes, preprint.
- [6] E. Savaş and S. Borgohain, On strongly almost lacunary statistical A -convergence defined by a Musielak-Orlicz function. (accepted in Filomat)

- [7] E. Savaş and S. Borgohain, Some new spaces of lacunary f -statistical A -convergent sequences of order α , *Advancements in Mathematical Sciences*, AIP Conf. Proc. 1676, 2015, 020086-1â€“020086-8; doi: 10.1063/1.4930512
- [8] E. Savaş, Generalized statistical convergence in random 2-normed space, *Iran. J. Sci. Technol. Trans. A Sci*, 36(2012), 417-423.
- [9] E. Savaş, I_λ -statistically convergent sequences in topological groups, *Acta Et Commentationes Universitatis Tartuensis De Mathematica*, 18(1)(2014), 33-38.
- [10] E. Savaş, On generalized sequence spaces via modulus function, *J. Inequal. Appl.*, 2014, 2014:101, 8 pp. 40A30.
- [11] E. Savaş, On some new sequence spaces defined by infinite matrix and modulus, *Adv. Difference Equ.*, 2013, 2013:274, 9 pp.
- [12] E. Savaş and R. F. Patterson, Double sequence spaces defined by a modulus. *Math. Slovaca*, 61(2), 245-256(2011).
- [13] E. Savaş and R. F. Patterson, Lacunary statistical convergence of multiple sequences, *Appl. Math. Lett.*, 19(6), 527-534(2006).
- [14] E. Savaş, On some generalized sequence spaces defined by a modulus, *Indian J. Pure Appl. Math.* 30(5), 459-464(1999).
- [15] E. Savaş, Strong almost convergence and almost λ -statistical convergence, *Hokkaido Math. J.*, 29(3), 531-536(2000).
- [16] H. Cakalli and E. Savaş, Statistical convergence of double sequences in topological groups, *J. Comput. Anal. Appl.*, 12(2), 421-426(2010).
- [17] H. I. Miller and C. Orhan, On almost convergent and statistically convergent subsequences, *Acta Math. Hungar.*, 93(1-2)(2001), 135-151.
- [18] H. Fast, Sur la convergence statistique, *Colloq. Math.* 2(1951), 241-244.
- [19] H. Nakano, Concave modulars, *J. Math. Soc. Japan.*, 5 (1953), 29â€“49.

- [20] I. Bala, V. K. Bhardwaj and E. Savaş, Some sequence spaces defined by $|A|$ summability and a modulus function in seminormed space, *Thai J. Math.*, 11(3), 623-631(2013).
- [21] I. J. Maddox , Sequence Spaces Defined by a modulus, *Mathematical Proceedings of the Cambridge Philosophical Society*, 100 (1986), 161-166.
- [22] I.J. Maddox, Inclusion between FK spaces and Kuttnerá's theorem, *Math. Proc. Camb. Philos. Soc.*, 101(1987), 523-527.
- [23] I.J. Schoenberg, The integrability of certain functions and related summability methods, *Amer. Math. Monthly*, 66(1959), 361-375.
- [24] J. Connor and E. Savaş, Lacunary statistical and sliding window convergence for measurable functions, *Acta Mathematica Hungarica*, 145(2)(2015), 416-432.
- [25] J.A. Fridy, On statistical convergence, *Analysis*, 5(1985), 301-313.
- [26] M. Mursaleen, λ -statistical convergence, *Math. Slovaca*, 50(2000), 111-115.
- [27] Richard F. Patterson and E. Savaş, Lacunary statistical convergence of double sequences, *Math. Commun.*, 10(1), 55-61(2005).
- [28] T. Šalát, On statistically convergent sequences of fuzzy real numbers, *Math. Slovaca*, 30(1980), 139-150.
- [29] W. H. Ruckle, FK spaces in which the sequence of coordinate vectors is bounded, *Canad. J. Math.*, 25 (1973), 973-978.